

Fitzroy estuary - macrobenthos

Fitzroy estuary assessment FE1 – Eco-health monitoring





Why use macrofauna to assess Fitzroy ecosystem health?

- Many organisms are relatively non-mobile and integrate effects of pollutants over time
- They are comparatively easy to quantitatively sample
- Major changes in keystone macrofaunal species can have profound impacts on higher trophic groups
- Facilitates comparative assessment of health in Port Curtis




Locations of Fitzroy River Macrobenthic Sampling Stations









Depth profiles for the Fitzroy River, Stations 1-8.






















⊨


Depth profiles for the Fitzroy River, Stations 9-16.




















Fitzroy River - schematic sampling design










Fitzroy River sediment profile









<u>Summary of taxonomic groups found in 15 of 80</u> <u>Fitzroy benthic grab samples</u>

Ē


Taxa	No. Species	No. Individuals
Polychaetes (worms)	15 (60%)	157 (7%)
Bivalves (shellfish)	3 (12%)	1989 (90%)
Crustaceans (crabs/prawns etc.)	7 (28%)	57 (3%)
Total	25	1989

Macro-invertebrates commonly encountered in the Fitzroy River







NIVIDS ordination of DSQRT Fitzroy estuary macrobenthos







 \equiv I otal sp. abundance superimposed on ordination of Fitzroy estuary macrobenthos



Shannon-Weiner diversity superimposed on ordination of Fitzroy estuary benthos

